

Drizzling Dithered WFPC2 Images—A Demonstration

Max Mutchler and Andrew Fruchter

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, Maryland
21218; e-mail mutchler@stsci.edu, fruchter@stsci.edu*

Abstract. The IRAF/STSDAS package **dither** and its primary task **drizzle** were created by Andrew Fruchter and Ivo Busko to reconstruct dithered WFPC2 images, and remove cosmic rays from singly-dithered images. Drizzling is akin to shifting-and-adding with a variable pixel size. The **drizzle** task was initially developed for the Hubble Deep Field project. Here, its use is demonstrated cookbook-style, using archival images of the edge-on galaxy NGC 4565 from Hubble Space Telescope (HST) program 6092.

1. Introduction

Although the HST and WFPC2 optics now provide an excellent PSF, the CCDs under-sample the images. On the three WF chips, the width of a pixel equals the FWHM of the PSF in the the near-infrared, and greatly exceeds it in the blue. The image quality can be improved by combining sub-pixel dithered images. If the dithers are particularly well-placed, the pixels from each image can simply be interlaced on a finer grid. But in practice, imperfect offsets and the geometric distortion can make interlacing impossible.

For purposes of combining the dithered images of the Hubble Deep Field, Richard Hook and Andrew Fruchter developed a new technique known as variable-pixel linear reconstruction, or “drizzling”. Drizzling can be thought of as a continuous set of linear functions that vary smoothly from the optimum linear combination technique—interlacing—to the old standby, shift-and-add. The degree to which one must depart from interlacing and move towards shift-and-add is determined by the nature of the input data. Drizzling naturally handles both missing data and geometric distortion, and can largely remove the effect on photometry produced by the geometric distortion of the WFPC2. For more background, see the articles listed in the reference section.

This paper shows how these tasks can be used to remove cosmic rays from singly-dithered images (i.e., in which only one exposure is taken at each dither position). The **dither** package in IRAF/STSDAS contains the following tasks:

```
precor      - Remove cosmic rays prior to cross-correlation
crossdriz   - Builds 1-group cross-correlation image (shift + rotation)
offsets     - Builds 4-group cross-correlation image (shift only)
shiftfind   - Finds x and y shifts in a cross-correlation image
avshift     - Averages the shifts measured on 4 WFPC chips
rotfind     - Finds rotation angle from a set of cross-correlation images
drizzle     - Perform linear image reconstruction
blot        - Inverse of drizzle
invert      - Inverts the weight masks
deriv       - Takes derivative of blotted images
drizcr2     - Combines cosmic ray masks and removes bad pixels
```




Figure 1. Mosaicked F814W image of NGC 4565 prior to drizzling—note the abundance of cosmic ray contamination. We will process only the WF2 chip in this example.

Note that the last three tasks listed above will not be available in IRAF/STSDAS until the next release. Also, the scripts `cosmic.calc` and `cosmic2.calc` must reside in your working directory for use by `drizcr2`. Scripts named `run*.cl` were used to ensure proper parameter settings at critical steps.

2. Measuring Shifts Between Dithered Images

The drizzling process is very sensitive to the accuracy of the shifts used, so we begin by carefully measuring the shifts between the dithered exposures. Our sample dataset is three singly-dithered WFPC2 images of the edge-on galaxy NGC 4565 from HST program 6092 by Keith Ashman (archival rootnames U31S0101T, U31S0102T, and U31S0103T). We will only work on the WF2 part of the image here.

2.1. Clean the Input Images

The measurement of shifts is often improved if some attempt to remove cosmic rays from the input images is made. Remove cosmic rays from the input images with `precor`. The clean images produced by `precor` should be used only for creating the cross-correlation images and measuring shifts, and NOT used as the input images in later stages.

```
di> precor *.hhh
```

2.2. Cross-correlate the Dithered Input Images

Create cross correlation images with the `crossdriz` task. We will use the first image as our reference image, and cross-correlate it with the 2nd and 3rd images:

```
cl> cl < runcrossdriz.cl
crossdriz.coeffs="drizzle$coeffs/trauger"
crossdriz.lambda=814
crossdriz ../102[2] ../101[2] 2x1
crossdriz ../103[2] ../101[2] 3x1
```

If we were measuring shifts for all four WFPC2 groups at the same time, the `offsets` task would be used instead of `crossdriz`.

2.3. Measure the Shifts Using the Cross-correlation Images

Use `shiftfind` to measure the shifts using the cross-correlation images (`*x1.hhh`). The output is written to a file named `shifts.out`. The shifts must be used consistently while running the `drizzle` and `blot` tasks later in this procedure:

```
cl> shiftfind *x1.hhh shifts.out
cl> more shifts.out
2x1.hhh    4.9714 0.0326    5.0206 0.0253
3x1.hhh   -5.0258 0.0330   -5.0148 0.0257
           x shift          y shift
```

In this example, we are measuring the shifts between WF2 images only. If we were measuring shifts on all four WFPC2 chips, we would run the `avshift` task at this point to average the shifts measured over all four chips:

```
cl> avshift shifts.out 0 > avshifts.out
```

3. Begin the Drizzling Procedure

Now that we have measured accurate shifts, we can begin drizzling. Create a subdirectory where we will work on one group (WF2 in this case) at a time. Then we `drizzle` the input images onto a finer grid, and unshift them. The output drizzled images are named `*p1.hhh` here to indicate that they were drizzled with `pixfrac=1.0` and to distinguish them from the final drizzled image (which will have `pixfrac=0.6` and be named `*p06.hhh`). Remember to `set stdimage=imt2048` to display the drizzled images on their new finer grid:

```
cl> cl < rundriz1.cl
drizzle.lambda=814
drizzle.wt_scl="exptime"
drizzle.expkey="exptime"
drizzle.pixfrac=1.0
drizzle.scale=0.5
drizzle.outnx=1600
drizzle.outny=1600
drizzle.coeffs=""
drizzle ../101.hhh[2] 101_p1 outweig=101_p1w xsh=0.0 ysh=0.0
drizzle ../102.hhh[2] 102_p1 outweig=102_p1w xsh=4.971 ysh=5.021
drizzle ../103.hhh[2] 103_p1 outweig=103_p1w xsh=-5.026 ysh=-5.015
```

4. Create a Median Image to Remove Cosmic Rays

Now we will combine the unshifted images from the previous step such that the median pixel value is used. The highest pixel values will be rejected, thereby removing cosmic rays.

4.1. Invert the Weight Images

Before we can combine the images, we must invert the output weighting images produced by `drizzle` (`*p1w.hhh`), since `gcombine` expects pixel values where good=0 and bad=1:

```
cl> invert
Mask(s) to be inverted: *p1w.hhh
```

4.2. Combine the Images

Use `gcombine` to create the median image from the WF2 I images (`wf2i_med.hhh`) using the inverted weight masks (`*p1w_inv.hhh`):

```
fo> gcombine *p1.hhh wf2i_med groups=1 masks=*p1w_inv.hhh
      reject=minmax combine=median weight=none scale=exposure
      expname=EXPTIME nlow=1 nhigh=1 rdnoise=5.2 gain=7.5
```

5. Blot the Median Image Back to the Input Plane

Use `blot` to make copies of the median image that are shifted back to the positions of the original input images. The output blotted images will be scaled to the input image exposure times and named `*bl.hhh`:

```
cl> cl < runblot.cl
blot.lambda=814
blot.scale=0.5
blot.outnx=800
blot.outny=800
blot.expout=600
blot.coeffs=""
blot wf2i_med 101_b1 xsh=0.0 ysh=0.0
blot wf2i_med 102_b1 xsh=4.971 ysh=5.021
blot wf2i_med 103_b1 xsh=-5.026 ysh=-5.015
```

5.1. Take the Derivative of the Blotted Images

Take the spatial derivative of each of the blotted images in preparation for the following step. Make sure you have loaded ALL the tasks called by `deriv`.

```
di> deriv
Image(s) for cosmic ray cleaning (@inlist-deriv): *_bl.hhh
```




Figure 2. A section of one of the three dithered F814W images of NGC 4565 with cosmic rays (above), and the final drizzled image with cosmic rays removed (below).

6. Combine Cosmic Ray Masks and Remove Bad Pixels

The `drizcr2` task compares the original images to their blotted counterparts. Any pixels that show a significant difference are flagged as bad pixels in the mask it creates. This task iterates to find bad pixels adjacent to others it has already found, since most cosmic ray hits involve many adjacent pixels. It uses the files `cosmic.calc` and `cosmic2.calc`, which must reside in your working directory. Compare the original images with their corresponding output masks to check that only truly bad pixels are being flagged:

```
cl> more cosmic.calc
if (abs(im1-im2) .gt. 0.4*im3+3.5*sqrt(7*abs(im2)+(7*2)*(7*2))/7) then 0 else 1

cl> more cosmic2.calc
if (im4 .lt. 9) then 0 else 1

cl> drizcr2
Image(s) for cosmic ray cleaning: ../10*.hhh
Group being cleaned: 2
```

7. Final Drizzling of the Input Images

Finally, re-drizzle the input images with the masks you have created, with a smaller `pixfrac` on a finer grid, with the geometric distortion coefficients applied:

```
cl> cl < rundriz2.cl
drizzle.lambda=814
drizzle.wt_scl="exptime"
drizzle.expkey="exptime"
drizzle.pixfrac=0.6
drizzle.scale=0.5
drizzle.coeffs="drizzle$coeffs/wf2-trauger"
drizzle.fillval=0
drizzle.outnx=1600
drizzle.outny=1600
drizzle ../101.hhh[2] wf2i_p06 outweig=wf2i_p06w in_mask=101_cr
xsh=0.0 ysh=0.0
drizzle ../102.hhh[2] wf2i_p06 outweig=wf2i_p06w in_mask=102_cr
xsh=4.971 ysh=5.021
drizzle ../103.hhh[2] wf2i_p06 outweig=wf2i_p06w in_mask=103_cr
xsh=-5.026 ysh=-5.015
```

The final output image is then `wf2i_p06.hhh`—the WF2 part of the F814W (I) image drizzled with a pixel fraction of 0.6. The process described here could now be repeated for each CCD and each color.

References

- Adorf, H., 1995, ST-ECF Newsletter, 23, 19
- Dickinson, M., & Fosbury, R., 1995, ST-ECF Newsletter, 22, 14
- Hook, R., & Fruchter, A., 1997, ST-ECF Newsletter, 24, 9
- Leitherer, C. (ed.), 1995, HST Data Handbook, version 2.0, Ch. 41, 520