



1255 - JWST Medium-Deep Fields -- Hammel IDS GTO Program

Cycle: 1, Proposal Category: GTO

INVESTIGATORS

<i>Name</i>	<i>Institution</i>	<i>E-Mail</i>
Dr. Stefanie N. Milam (PI)	NASA Goddard Space Flight Center	stefanie.n.milam@nasa.gov
Dr. Rolf A. Jansen (CoI)	Arizona State University	rolf.jansen@asu.edu
Dr. Heidi B. Hammel (CoI)	Space Science Institute	hbhammel@aura-astronomy.org

OBSERVATIONS

<i>Folder</i>	<i>Observation</i>	<i>Label</i>	<i>Observing Template</i>	<i>Science Target</i>
NEP Time-Domain Field				
	1	NEP TDS FIELD Spoke 5	NIRCam Imaging	(1) JWST-NEP-TDS-FIELD

ABSTRACT

Our team assessed the detectability of distant solar system objects based on the parameters discussed in the “JWSTNEPcommunityfield16” document provided by JWST IDS R. Windhorst. Overall, nearly all distant solar system populations (> 40 AU) would have very low detectability: the proposed field is near the ecliptic pole (generally rules out Kuiper Belt Objects both classical and scattered, as well as Inner Oort Cloud Objects); the objects are extremely faint (e.g., Oort Cloud Objects); or the object's low density across the sky precludes meaningful discovery statistics (the near-polar-orbit Centaurs). There may be some populations that we know very little about such as bodies on near-polar orbits that could be detectable, but their numbers are not guaranteed. We nonetheless felt that given the non-zero probability of detection and given the importance of an actual detection, it was worthwhile to contribute time to the survey.

This includes observation IDs: Hammel_6000, 6001, 6002, 6003, 6004, 6005, 6006, 6007, 6008, 6009, 6010.

OBSERVING DESCRIPTION

JWST NEP Time-Domain Community Field: Our team assessed the detectability of distant solar system objects based on the parameters discussed in the “JWSTNEPcommunityfield16” document provided by JWST IDS R. Windhorst. Overall, nearly all distant solar system populations (> 40 AU) would have very low detectability: the proposed field is near the ecliptic pole (generally rules out Kuiper Belt Objects both classical and scattered, as well as Inner Oort Cloud Objects); the objects are extremely faint (e.g., Oort Cloud Objects); or the object's low density across the sky precludes meaningful discovery statistics (the near-polar-orbit Centaurs). There may be some populations that we know very little about such as bodies on near-polar orbits that could be detectable, but their numbers are not guaranteed. We nonetheless felt that given the non-zero probability of detection and given the importance of an actual detection, it was worthwhile to contribute time to the survey.

Within the northern CVZ, in the very best field for Time-Domain Science with JWST, IDS Windhorst defined a ~46 hr Medium-Deep Blank Field GTO program, covering a 4-spoke pattern that can efficiently provide NIRISS slitless grism spectroscopy of a large portion of the area observed with NIRCcam. Each spoke in this unique pattern consists of a 1row x 2col NIRCcam mosaic with significant overlap to fill the large (~44") gap between the NIRCcam modules and to provide contiguous NIRISS spectroscopic coverage. At each mosaic pointing, we fill the 4"-5" intra-module gaps using a 3-step INTRAMODULE dither (no sub-pixel dithering).

The present observations add half of a component for an additional spoke at an orientation that notionally differs by 45 degrees from the pattern defined by Windhorst to add both areal coverage over which full object characterization is obtained, as well as time-domain coverage in the areas (wedges) of overlap. NIRCcam imaging is obtained in F090W, F115W, F150W, F200W in the SW, and F277W, F356W, F444W, and F410M in the LW channel, with the F150W, F200W, F277W and F356W exposures split to match number of NIRISS exposures). In parallel to the NIRCcam imaging, we obtain NIRISS slitless spectroscopy with the orthogonal F150C and F150R grisms, crossed with the F200W filter (the associated NIRISS direct images are with the F200W filter).

Proposal 1255 - Targets - JWST Medium-Deep Fields -- Hammel IDS GTO Program

Fixed Targets	#	Name	Target Coordinates	Targ. Coord. Corrections	Miscellaneous
	(1)	JWST-NEP-TDS-FIELD	RA: 17 22 47.8960 (260.6995667d) Dec: +65 49 21.54 (65.82265d) Equinox: J2000 <i>Comments: PA offset from WINDHORST_1131 by 255 to 315 degrees</i> <i>Category=Unidentified</i> <i>Description=[Infrared sources, Radio sources, Variable radiation sources, Visible sources, X-ray sources]</i> <i>Extended=NO</i>		

Proposal 1255 - Observation 1 - JWST Medium-Deep Fields -- Hammel IDS GTO Program

Tue Feb 05 22:33:26 GMT 2019

Observation	Proposal 1255, Observation 1: NEP TDS FIELD Spoke 5 Diagnostic Status: Warning Observing Template: NIRCam Imaging Coordinated Parallel Template(s): NIRISS Wide Field Slitless Spectroscopy										
	(Visit 1:1) Warning (Form): Overheads are provisional until the Visit Planner has been run.										
Fixed Targets	#	Name	Target Coordinates			Targ. Coord. Corrections			Miscellaneous		
	(1)	JWST-NEP-TDS-FIELD	RA: 17 22 47.8960 (260.6995667d) Dec: +65 49 21.54 (65.82265d) Equinox: J2000								
<i>Comments: PA offset from WINDHORST_1131 by 255 to 315 degrees</i> <i>Category=Unidentified</i> <i>Description=[Infrared sources, Radio sources, Variable radiation sources, Visible sources, X-ray sources]</i> <i>Extended=NO</i>											
Template	NIRCam Imaging					NIRISS Wide Field Slitless Spectroscopy					
	Module: ALL Subarray: FULL										
Dithers	#	Primary Dither Type		Primary Dithers	Dither Size	Subpixel Positions		Coordinated Parallel Subpixel Selector		Dither Direct Images Primes	
	1	INTRAMODULE		3		1		NIRCam Only		NO_DITHERING	
Spectral Elements	NIRCam Imaging	Short Filter	Long Filter	Readout Pattern	Groups/Int	Integrations/Exp	Total Integrations	Total Dithers	Total Exposure Time	ETC Wkbk.Calc ID	
	1	F150W	F356W	SHALLOW2	4	2	2	1	386.524		
	2	F090W	F444W	BRIGHT2	3	1	3	3	225.472		
	3	F150W	F356W	SHALLOW2	4	2	2	1	386.524		
	4	F200W	F277W	SHALLOW2	4	2	2	1	386.524		
	5	F115W	F410M	BRIGHT2	3	1	3	3	225.472		
	6	F200W	F277W	SHALLOW2	4	2	2	1	386.524		
Spectral Elements	NIRISS Wide Field Slitless Spectroscopy	Exposure Type	Filter	Grism	Readout Pattern	Groups/Int	Integrations/Exp	Total Dithers	Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
	1	DIRECT	F200W		NIS	9	1	1	1	397.26	
	2	GRISM	F200W	GR150C	NIS	2	1	3	3	289.893	
	3	DIRECT	F200W		NIS	9	1	1	1	397.26	
	4	DIRECT	F200W		NIS	9	1	1	1	397.26	
	5	GRISM	F200W	GR150R	NIS	2	1	3	3	289.893	
	6	DIRECT	F200W		NIS	9	1	1	1	397.26	

Proposal 1255 - Observation 1 - JWST Medium-Deep Fields -- Hammel IDS GTO Program

Special Requirements

Aperture PA Range 110 to 120 Degrees (V3 110.112526 to 120.112526)
Offset 190.0 arcsec, -105.0 arcsec
No Parallel