A Dedicated Space Telescope for Time-Domain Solar System Science

Michael H. Wong
mikewong@astro.berkeley.edu
STScI (Instruments Division)
UC Berkeley (Astronomy Department)

Máté Ádámkovics
UC Berkeley

Sushil K. Atreya
Univ. Mich.

Don Banfield
Cornell

Jim Bell
Cornell

Susan Benecchi
STScI

Gordon Bjoraker
NASA GSFC

John R. Casani
JPL

John T. Clarke
Boston Univ.

Imke de Pater
UC Berkeley

Scott G. Edgington
JPL

Leigh N. Fletcher
JPL

Richard G. French
Wellesley

William Grundy
Lowell

Amanda R. Hendrix
JPL

Daniel Hestroffer
Obs. Paris Meudon

Erich Karkoschka
Univ. Arizona

Jian-Yang Li
Univ. Maryland

Franck Marchis
SETI/UC Berkeley

Melissa A. McGrath
NASA MSFC

William J. Merline
SwRI

Julianne I. Moses
LPI

Keith Noll
STScI

Glenn S. Orton
JPL

Kathy A. Rages
SETI

Kurt Retherford
SWRI

Joachim Saur
Univ. Cologne

Kunio Sayanagi
Caltech

Nick Schneider
CU

Eric H. Smith
Lockheed Martin

Lawrence A. Sromovsky
UW-Madison

Nathan J. Strange
JPL

Anne Verbiscer
Univ. Virginia

Padmavati A. Yanamandra-Fisher
JPL

Abstract

Time-variable phenomena with scales ranging from minutes to decades have led to a large fraction of recent advances in many aspects of solar system science. ***We present the scientific motivation for a dedicated space observatory for solar system science.*** We call the observatory the Planetary Dynamics Explorer (PDX). This facility will ideally conduct repeated imaging and spectroscopic observations over a period of 10 years or more. It will execute a selection of long-term projects with interleaved scheduling, resulting in the acquisition of data sets with consistent calibration, long baselines, and optimized sampling intervals.

A sparse aperture telescope would be an ideal configuration for the mission, trading decreased sensitivity for reduced payload mass, while preserving spatial resolution. Ultraviolet capability is essential, especially once the Hubble Space Telescope retires.

Specific investigations will include volcanism and cryovolcanism (on targets such as Io, Titan, Venus, Mars, and Enceladus); seasonal weather cycles; zonal flow, vortices, and storm evolution on the giant planets; mutual events and orbit determination of multiple small solar system bodies; auroral, solar wind, and magnetospheric interactions; variability in thin atmospheres; and cometary evolution. The mission will produce a wealth of data products—such as long time-lapse movies of planetary atmospheres—with outreach potential as well as scientific value.

Existing and planned ground- and space-based facilities are not suitable for these time-domain optimized planetary dynamics studies for numerous reasons, including: oversubscription by astrophysical users, field-of-regard limitations, sensitive detector saturation limits that preclude bright planetary targets, and limited mission duration.

Requirements for PDX

The chief requirements for time-domain solar system science are angular resolution, sampling interval, and campaign duration. Although many ground- and space-based telescopes satisfy the angular resolution requirement, only a dedicated solar system mission could achieve the time domain requirements.

Angular resolution

Studies of planetary dynamics require the resolution of small distant objects such as cloud features, volcanic plumes, and fragments of small solar system bodies. With a nominal 3-m aperture, PDX will achieve an angular resolution of about 40 mas at optical wavelengths. This resolution is comparable to that provided by HST and the best current ground-based telescopes, which have demonstrated a wealth of time-domain science opportunities (see below). In the coming decades, extremely high resolution will be afforded by very large ground-based telescopes, but PDX will not attempt to compete with those efforts, instead specifying a 40 mas resolution based on the minimum requirement to image dynamically relevant features in the solar system.

Sampling interval

Observing programs will be scheduled to ensure that each program acquires data at its desired sampling interval, which will typically range from hours to days. Occultation light curves will push the short-interval limits with millisecond-range sampling intervals.

Campaign duration

Campaign durations lasting from the entire mission lifetime to single visits will be accommodated, providing new opportunities as compared with semester- or cycle-based scheduling at other observatories. In particular, campaigns lasting the full mission lifetime will enable high-return high-risk science such as cryovolcanic activity searches.

Resolution / sensitivity / volume trades

For most space telescope missions, photometric sensitivity is a major requirement. For time-domain solar system astronomy, a trade can be studied between sensitivity and payload volume, since relatively bright solar system targets are less demanding of sensitivity. A sparse-aperture configuration enables this trade to favor smaller payload volume, maximizing the total aperture (and thus resolving power) for a given payload size.



Figure 1. Sparse aperture telescope examples. *Top left:* Illustration of the MIDAS multiple telescope array for spaceborne remote sensing (Smith et al. 2005). *Middle and bottom left:* The Star-9 prototype distributed aperture telescope on its optical testbed at Lockheed Martin (Rieboldt et al. 2005, Kendrick et al. 2006). *Right:* Optical path of the Star-9 system (Kendrick et al. 2005). ©2006 Lockheed Martin Corporation. Images from Kendrick et al. (2006), courtesy of Lockheed Martin Corporation.

Science programs

PDX will provide an ideal platform for a range of time-domain research programs. A selection of general observer programs will be solicited from the broader solar system research community. Programs will be prioritized based on their scientific impact derived from the effective use of the resolution, sampling rate, and campaign duration opportunities uniquely provided by the PDX observatory. A set of observing programs with complementary temporal requirements will be chosen to maximize the facility duty cycle. Monitoring programs will be able to trigger shorter-interval target of opportunity observations under the appropriate conditions.

PDX will make its greatest contributions in the areas of planetary atmospheres, active geology, and small solar system bodies. A sample of candidate investigations is given in this section; many more investigations could be pursued, but these give at least a sense of the demands on sampling interval and campaign duration that are important for time-domain solar system science.

Table 1. Examples of time-domain solar system science investigations that could be explored using PDX. Note that since most investigations requiring spectroscopic data also require spatially resolved spectra, an integral field spectrometer might be an ideal spectroscopic instrument choice.

Investigation	Category	Data type (wavelength regime)	Sampling scales	Campaign duration
Atmopheric dynamics: winds and transport	Atmospheres	Imaging (O)	Hours, days	Years
Variable atmos. structure & composition	Atmospheres	Imaging, spectroscopy (UV, O, IR)	Hours, days	Days, years
Occultations	Atmospheres	Photometry, spectroscopy (UV, O, IR)	Milliseconds	Hours
Aurorae, magnetospheres	Atmospheres/space science	Imaging, spectroscopy (UV)	Minutes, hours	Years, hours
Volcanic trace gases	Atmospheres/geology/astrobiology	Spectroscopy, imaging (IR)	Days	Years
Volcanic plumes	Geology	Imaging, spectroscopy (O, IR)	Days, hours	Years
Cryovolcanism	Geology/astrobiology	Imaging, spectroscopy (UV, O, IR)	Days	Years
Mutual events, lightcurves	Small bodies	Photometry (O)	Milliseconds, minutes	Hours, months
Cometary evolution	Small bodies	Imaging, spectroscopy (UV, O, IR)	Hours	Days



Figure 2. Examples of time-domain science at Jupiter. HST ACS/HRC images of Jupiter from 2006 at ~50 mas effective resolution *(A)* were used to provide velocity profiles along the east-west *(B)* and north-south *(C)* axes of Oval BA (Asay-Davis et al. 2009). Differences between 2000 and 2006 velocity profiles near 30° south latitude are attributed to transient interactions (of unknown timescale) between the vortex and its northward jet. Wind measurements require sampling intervals on hour timescales. Zonal wind profiles before *(D; black)*, during *(D; grey)*, and after *(E)* the eruptions of massive powerful convective plumes *(G)* revealed significant changes in the zonal winds *(F)* as derived from HST WFPC2 imaging data (Sánchez-Lavega et al. 2008).



Figure 3. *Left:* Spectral imaging observations of TItan using VLT/SINFONI with ~45 mas effective resolution show that observations at 1-day intervals can reveal significant cloud variation. Much higher spatial resolution Cassini ISS data showed even more rapid change at hour timescales, but it is not clear that PDX could spatially resolve the rapid changes. *Below:* Schaller et al. (2006) used observations taken at day-scale intervals to demonstrate changes in cloud activity, which they relate to a large convective event in 2004 and to seasonal effects. The 2002 vertical line is southern solstice; the 2005 vertical line is the date when Titan's region of maximum insolation moved north from the south pole.





Figure 4. Images of Io at ~50 mas resolution demonstrate that a sampling interval of no greater than 1 day is probably required to catch the development of volcanic eruptions (here, the volcano Surt; see Marchis et al. 2002). However, eruption durations last for days to months, so sampling intervals could be tuned depending on specific science goals.



Figure 5. Although taken at different wavelengths, these 1986 Voyager *(top)* and 2005–2007 Keck (*bottom)* images of Uranus demonstrate the variability of cloud activity over long timescales. The bright cloud complex (bottom left) has been tracked for several years and changes in altitude, area, latitude, and longitude (Sromovsky et al. 2007). Observations spanning many years—including data from HST/WFPC2—show evidence of significant decade-scale changes on both Uranus and Neptune that is only partially explicable as effects of changing subsolar latitude (Rages et al. 2004, Hammel and Lockwood 2007).

References

Asay-Davis, X., Marcus, P.S., Wong, M.H., de Pater, I., 2009. Jupiter's evolving Great Red Spot: Velocity measurements with the ACCIV automated cloud tracking method. *Icarus*, in press.

Hammel, H.B., Lockwood, G.W., 2007. Long-term atmospheric variability on Uranus and Neptune. *Icarus* 186, 291–301.

Kendrick, R.L., and 40 authors, 2006. Wide-field Fizeau imaging telescope: experimental results. Applied Optics 45, 4235–4240.

Marchis, F., de Pater, I., Davies, A.G., Roe, H.G., Fusco, T., Le Mignant, D., Descamps, P., Macintosh, B.A., Prangé, R., 2002. High-Resolution Keck Adaptive Optics Imaging of Violent Volcanic Activity on Io. *Icarus* 160, 124–131.

Rages, K.A., Hammel, H.B., Friedson, A.J., 2004. Evidence for temporal change at Uranus' south pole. *Icarus* 172, 548–554.

Rieboldt, S.E., Wong, M.H., Ádámkovics, M., Delory, G.T., de Pater, I., Manga, M., Lipps, J.H., Dalton, J.B., Pitman, J., Kendrick, R.L., 2005. MIDAS: Advanced Remote Sensing for the Exploration of Icy Satellites. American Geophysical Union, Fall Meeting 2005, abstract #P11B-0121.

Sánchez-Lavega, A., G.S. Orton, R. Hueso, E. García-Melendo, S. Pérez-Hoyos, A. Simon-Miller, J.F. Rojas, J.M. Gómez, P. Yanamandra-Fisher, L. Fletcher, J. Joels, J. Kemerer, J. Hora, E. Karkoschka, I. de Pater, M.H. Wong, P.S. Marcus, N. Pinilla-Alonso, and the IOPW team, 2008. Depth of a strong jovian jet from a planetary-scale disturbance driven by storms. *Nature* 451, 437–440.

Schaller, E.L., Brown, M.E., Roe, H.G., Bouchez, A.H., Trujillo, C.A., 2006. Dissipation of Titan's south polar clouds. *Icarus* 184, 517–523.

Smith, E.H., de Leon, E., Dean, P., Deloumi, J., Duncan, A., Hoskins, W., Kendrick, R., Mason, J., Page, J., Phenis, A., Pitman, J., Pope, C., Privari, B., Ratto, D., Romero, E., Shu, K.-L., Sigler, R., Stubbs, D., Tapos, F., Yee, A., 2005. Multiple instrument distributed aperture sensor (MIDAS) testbed. *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series* 5882, 488–500.

Sromovsky, L.A., Fry, P.M., Hammel, H.B., de Pater, I., Rages, K.A., Showalter, M.R., 2007. Dynamics, evolution, and structure of Uranus' brightest cloud feature. *Icarus* 192, 558–575.